FICHA TÉCNICA

Línea: MELINE®

Producto: 02 MELINE® DARK CIRCLES



Descripción e indicación: Producto de aplicación domiciliaria para la mejora y optimización de los resultados obtenidos con 01 MELINE® Dark Circles.

Mecanismo de acción: Combinación de activos despigmentantes dermocosmeticos que han sido seleccionados por su alta tolerancia cutánea en la zona periocular. Actúa mejorando y atenuando las ojeras, disminuyendo los depósitos de melanina formada, así como las pigmentaciones de origen melanohemático. Además, estimula el ciclo celular luchando contra los signos visibles del fotoenvejecimiento de la piel.

Modo de Uso: Aplicar mañana y noche, con la ayuda del roll-on, una capa sobre la piel de la zona periocular y dejar secar. Utilizar durante un tiempo mínimo no inferior a 30 días, empezando su uso el 2º día tras la aplicación de 01 MELINE® Dark Circles.

Principios activos y propiedades:

- Ruscus Aculeatus, Gingko Biloba,Troxerutina y Melilotus Officinalis: Activos venotónicos y linfoquinéticos que permiten una acción descongestionante sobre el área periocular.

- Alfa Arbutina y Ácido Kójico: Acción inhibitoria de la Tirosinasa.

- Acido Tranexámico: Disminuye la actividad inflamatoria y vascular relacionada con la pigmentación periocular.

- Matrikinas y Flavonoides: Refuerzan la firmeza y el tono del contorno del ojo, además de contribuir a la eliminación de pigmentos causantes del oscurecimiento periocular.

- Vitamina P (flavonoides, saponósidos y procianidoles): favorecen la microcirculación y el fortalecimiento de las paredes venosas.

Presentación: 10 ml.

Bibliografía:

- *Stern EC. Topical application of lactic acid in the treatment and prevention of certain disorders of the skin. Urol Cutaneous Rev 1946;50:106-7.*
- *Chiu A, Kimball AB. Topical vitamins: Mineral and botanical ingredients as modulators of environmental and chronological skin damage. Br J Dermatol 2003;149:681-91.*
- *Elson M, Nacht S. Treatment of periorbital hyperpigmentation with topical vitamin K/vitamin A. J Cosmetic Dermatol 1999;12:323-5.*
- *Kaczvinsky JR, Griffiths CE, Schnicker MS, Li J. Efficacy of anti-aging products for periorbital wrinkles as measured by 3-D imaging. J Cosmet Dermatol 2009;8:228-33.*
- *Edison BL, Green BA, Wildnauer RH, Sigler ML. A polyhydroxy acid skin care regimen provides anti-aging effects comparable to an alpha-hydroxy acid regimen. Cutis 2004; 73:14-7.*
- *Kim EH, Kim YC, Lee ES, Kang HY. The vascular characteristics of melasma. J Dermatol Sci 2007; 46: 111–16.*
- *Noh TK, Choi SJ, Chung BY et al. Inflammatory features of melasma lesions in Asian skin. J Dermatol 2014; 41: 788–94.*
- *Tse TW, Hui E. Tranexamic acid: an important adjuvant in the treatment of melasma. J Cosmet Dermatol 2013; 12: 57–66.*
- *Steiner D, Feola C, Bialeski N et al. Tranexamic acid in treatment of melasma. Surg Cosmetic Dermatol 2009; 1: 174–7.*
- *Bunney, S. 1992. The Illustratcd Encyclopedia of Herb. Their Medicinal and Culinary uses, Chanceller Press London..190.*
- *lkuta, A. and Itokawa, H. 1988. Triterpenoids of Paeonia Japonica callus tissue. Phytochemistry 27: 2813.*
- *Markham, KR. and Ternai, B. 1976. Tetrahedron, 32: 2607.*
- *Nagar, A., Gujral, VK. and Gupta, BA. 1979. Arjunone, A new flavanone from T. arjuna, its constitution and synthesis. Planta Medica. 37: 183.*

***